

---

# Nucleic Acids Research

---

## Contents

Volume 35 number 2

---

### CHEMISTRY

Efficient signaling platforms built from a small catalytic DNA and doubly labeled fluorogenic substrates

W.Chiuman and Y.Li

401–405

**S** Antisense oligonucleotides containing locked nucleic acid improve potency but cause significant hepatotoxicity in animals

E.E.Swayze, A.M.Siwkowski, E.V.Wancewicz, M.T.Migawa, T.K.Wyrzykiewicz, G.Hung, B.P.Monia and C.F.Bennett

687–700

### COMPUTATIONAL BIOLOGY

**S** More complete gene silencing by fewer siRNAs: transparent optimized design and biophysical signature

I.Ladunga

433–440

**M, S** High sensitivity RNA pseudoknot prediction

X.Huang and H.Ali

656–663

**M, S** Dynamic use of multiple parameter sets in sequence alignment

X.Huang and D.L.Brutlag

678–686

### GENOMICS

**M** G-quadruplexes in promoters throughout the human genome

J.L.Huppert and S.Balasubramanian

406–413

**M** GISMO—gene identification using a support vector machine for ORF classification

L.Krause, A.C.McHardy, T.W.Nattkemper, A.Pühler, J.Stoye and F.Meyer

540–549

**M** Genome-wide analyses of retrogenes derived from the human box H/ACA snoRNAs

Y.Luo and S.Li

559–571

### MOLECULAR BIOLOGY

**S** Role of Elg1 protein in double strand break repair

H.Ogiwara, A.Ui, T.Enomoto and M.Seki

353–362

**S** Improved nuclear localization of DNA-binding polyamides

N.G.Nickols, C.S.Jacobs, M.E.Farkas and P.B.Dervan

363–370

**S** SECIS elements in the coding regions of selenoprotein transcripts are functional in higher eukaryotes

H.Mix, A.V.Lobanov and V.N.Gladyshev

414–423

**S** Chloroplast phosphoglycerate kinase, a gluconeogenic enzyme, is required for efficient accumulation of *Bamboo mosaic virus*

J.-W.Lin, M.-P.Ding, Y.-H.Hsu and C.-H.Tsai

424–432

Ascorbate acts as a highly potent inducer of chromate mutagenesis and clastogenesis: linkage to DNA breaks in G<sub>2</sub> phase by mismatch repair

M.Reynolds, L.Stoddard, I.Bespalov and A.Zhitkovich

465–476

Glucocorticoid-induced leucine zipper (GILZ)/NF- $\kappa$ B interaction: role of GILZ homo-dimerization and C-terminal domain

B.D.Marco, M.Masseti, S.Bruscoli, A.Macchiarulo, R.Di.Virgilio, E.Velardi, V.Donato, G.Migliorati and C.Riccardi

517–528

*Continued*

	Two primate-specific small non-protein-coding RNAs in transgenic mice: neuronal expression, subcellular localization and binding partners	T.Khanam, T.S.Rozhdestvensky, M.Bundman, C.R.Galiveti, S.Handel, V.Sukonina, U.Jordan, J.Brosius and B.V.Skryabin	529–539
<b>S</b>	Expression of RAB4B, a protein governing endocytic recycling, is co-regulated with MHC class II genes	M.Krawczyk, E.Leimgruber, Q.Seguín-Estévez, I.Dunand-Sauthier, E.Barras and W.Reith	595–605
	Analysis of the DNA-binding sequence specificity of the archaeal transcriptional regulator Ss-LrpB from <i>Sulfolobus solfataricus</i> by systematic mutagenesis and high resolution contact probing	E.Peeters, C.Wartel, D.Maes and D.Charlier	623–633
<b>S</b>	A novel E4BP4 element drives circadian expression of <i>mPeriod2</i>	T.Ohno, Y.Onishi and N.Ishida	648–655
<b>S</b>	The bidirectional promoter of two genes for the mitochondrial translational apparatus in mouse is regulated by an array of CCAAT boxes interacting with the transcription factor NF-Y	E.Zanotto, Z.H.Shah and H.T.Jacobs	664–677
<b>NUCLEIC ACID ENZYMES</b>			
<b>S</b>	Association of Dnmt3a and thymine DNA glycosylase links DNA methylation with base-excision repair	Y.-Q.Li, P.-Z.Zhou, X.-D.Zheng, C.P.Walsh and G.-L.Xu	390–400
<b>S</b>	Nucleus-encoded plastid sigma factor SIG3 transcribes specifically the <i>psbN</i> gene in plastids	W.Zghidi, L.Merendino, A.Cottet, R.Mache and S.Lerbs-Mache	455–464
<b>S</b>	Flexible segments modulate co-folding of dUTPase and nucleocapsid proteins	V.Németh-Pongrácz, O.Barabás, M.Fuxreiter, I.Simon, I.Pichová, M.Rumlová, H.Zábranská, D.Svergun, M.Petoukhov, V.Harmat, É.Klement, É.Hunyadi-Gulyás, K.F.Medzihradzky, E.Kónya and B.G.Vértessy	495–505
<b>S</b>	Functional organization of the Rpb5 subunit shared by the three yeast RNA polymerases	C.Zaros, J.-F.Briand, Y.Boulard, S.Labarre-Mariotte, M.C.Garcia-Lopez, P.Thuriaux and F.Navarro	634–647
<b>RETRACTION</b>			
	Non-random, individual-specific methylation profiles are present at the sixth CTCF binding site in the human H19/IGF2 imprinting control region	J.Tost, H.Jammes, J.-M.Dupont, C.Buffat, B.Robert, T.-M.Mignot, F.Mondon, B.Carbonne, U.Siméoni, G.Grangé, A.Kerjean, F.Ferré, I.G.Gut and D.Vaiman	701
<b>RNA</b>			
<b>S</b>	Cold shock domain proteins and glycine-rich RNA-binding proteins from <i>Arabidopsis thaliana</i> can promote the cold adaptation process in <i>Escherichia coli</i>	J.S.Kim, S.J.Park, K.J.Kwak, Y.O.Kim, J.Y.Kim, J.Song, B.Jang, C.-H.Jung and H.Kang	506–516
	Incorporation of 5-fluorouracil into U2 snRNA blocks pseudouridylation and pre-mRNA splicing <i>in vivo</i>	X.Zhao and Y.-T.Yu	550–558
	Interplay of ‘induced fit’ and preorganization in the ligand induced folding of the aptamer domain of the guanine binding riboswitch	J.Noeske, J.Buck, B.Fürtig, H.R.Nasiri, H.Schwalbe and J.Wöhnert	572–583
	Reduced splicing efficiency induced by synonymous substitutions may generate a substrate for natural selection of new splicing isoforms: the case of CFTR exon 12	M.Raponi, F.E.Baralle and F.Pagani	606–613
	Encoding folding paths of RNA switches	A.Xayaphoummine, V.Viasnoff, S.Harlepp and H.Isambert	614–622

## STRUCTURAL BIOLOGY

	<b>Modulating role of RNA structure in alternative splicing of a critical exon in the spinal muscular atrophy genes</b>	N.N.Singh, R.N.Singh and E.J.Androphy	371–389
	<b>L11 domain rearrangement upon binding to RNA and thiostrepton studied by NMR spectroscopy</b>	H.R.A.Jonker, S.Ilin, S.K.Grimm, J.Wöhnert and H.Schwalbe	441–454
	<b>Selenium derivatization of nucleic acids for crystallography</b>	J.Jiang, J.Sheng, N.Carrasco and Z.Huang	477–485
<b>S</b>	<b>Influence of cationic molecules on the hairpin to duplex equilibria of self-complementary DNA and RNA oligonucleotides</b>	S-i.Nakano, T.Kirihata, S.Fujii, H.Sakai, M.Kuwahara, H.Sawai and N.Sugimoto	486–494
	<b>Structural basis for sequence-dependent DNA cleavage by nonspecific endonucleases</b>	Y.-T.Wang, W.-J.Yang, C.-L.Li, L.G.Doudeva and H.S.Yuan	584–594

## METHODS

<b>S</b>	<b>Positional artifacts in microarrays: experimental verification and construction of COP, an automated detection tool</b>	H.Yu, K.Nguyen, T.Royce, J.Qian, K.Nelson, M.Snyder and M.Gerstein	e8 (12 pp.)
<b>S</b>	<b>Human <i>RAD18</i> is involved in S phase-specific single-strand break repair without PCNA monoubiquitination</b>	N.Shiomi, M.Mori, H.Tsuji, T.Imai, H.Inoue, S.Tateishi, M.Yamaizumi and T.Shiomi	e9 (12 pp.)
	<b>Development of a species-specific RNA polymerase I-based shRNA expression vector</b>	M.S.Brenz Verca, P.Weber, C.Mayer, C.Graf, D.Refojo, R.Kühn, I.Grümmt and B.Lutz	e10 (8 pp.)
	<b>A semi-automated high-throughput approach to the generation of transposon insertion mutants in the nematode <i>Caenorhabditis elegans</i></b>	Y.Duverger, J.Belougne, S.Scaglione, D.Brandli, C.Beclin and J.J.Ewbank	e11 (8 pp.)
<b>S</b>	<b>A pHMM-ANN based discriminative approach to promoter identification in prokaryote genomic contexts</b>	S.Mann, J.Li and Y.-P.P.Chen	e12 (7 pp.)
	<b>Increasing the sensitivity of DNA microarrays by metal-enhanced fluorescence using surface-bound silver nanoparticles</b>	C.R.Sabanayagam and J.R.Lakowicz	e13 (9 pp.)

## ERRATUM

704

## CORRIGENDUM

702

703

*M*, contains a novel method

*S*, Supplementary Material available at NAR Online