
Nucleic Acids Research

Contents

Volume 35 number 9

CHEMISTRY

- Investigation of spectral conversion of d(TTAGGG)₄ and d(TTAGGG)₁₃ upon potassium titration by a G-quadruplex recognizer BMVC molecule C.-C.Chang, C.-W.Chien, Y.-H.Lin, C.-C.Kang and T.-C.Chang 2846–2860

COMPUTATIONAL BIOLOGY

- M* RNAmmer: consistent and rapid annotation of ribosomal RNA genes K.Lagesen, P.Hallin, E.A.Rødland, H.-H.Stærfeldt, T.Rognes and D.W.Ussery 3100–3108

GENOMICS

- S* Mapping translocation breakpoints using a wheat microarray P.R.Bhat, A.Lukaszewski, X.Cui, J.Xu, J.T.Svensson, S.Wanamaker, J.G.Waines and T.J.Close 2936–2943
- S* Sequence analysis of p53 response-elements suggests multiple binding modes of the p53 tetramer to DNA targets B.Ma, Y.Pan, J.Zheng, A.J.Levine and R.Nussinov 2986–3001
- S* Mitochondrial diversity within modern human populations R.W.Carter 3039–3045

MOLECULAR BIOLOGY

- M* Recovery of bisulfite-converted genomic sequences in the methylation-sensitive QPCR K.Munson, J.Clark, K.Lamparska-Kupsik and S.S.Smith 2893–2903
- S* Recruitment of DNA repair synthesis machinery to sites of DNA damage/repair in living human cells K.Hashiguchi, Y.Matsumoto and A.Yasui 2913–2923
- DNA modification by sulfur: analysis of the sequence recognition specificity surrounding the modification sites J.Liang, Z.Wang, X.He, J.Li, X.Zhou and Z.Deng 2944–2954
- All APOBEC3 family proteins differentially inhibit LINE-1 retrotransposition M.Kinomoto, T.Kanno, M.Shimura, Y.Ishizaka, A.Kojima, T.Kurata, T.Sata and K.Tokunaga 2955–2964
- Complex translational regulation of BACE1 involves upstream AUGs and stimulatory elements within the 5' untranslated region M.Mihailovich, R.Thermann, F.Grohovaz, M.W.Hentze and D.Zacchetti 2975–2985
- RNA polymerase III transcription is repressed in response to the tumour suppressor ARF J.P.Morton, T.Kantidakis and R.J.White 3046–3052
- Mechanism for the TtDnaA–Tt-oriC cooperative interaction at high temperature and duplex opening at an unusual AT-rich region in *Thermoanaerobacter tengcongensis* H.Pei, J.Liu, J.Li, A.Guo, J.Zhou and H.Xiang 3087–3099
- S* Actin-related protein Arp4 functions in kinetochore assembly H.Ogiwara, A.Ui, S.Kawashima, K.Kugou, F.Onoda, H.Iwahashi, M.Harata, K.Ohta, T.Enomoto and M.Seki 3109–3117

Continued

NUCLEIC ACID ENZYMES

- S* The unstructured C-terminus of the τ subunit of *Escherichia coli* DNA polymerase III holoenzyme is the site of interaction with the α subunit S.Jergic, K.Ozawa, N.K.Williams, X.-C.Su, D.D.Scott, S.M.Hamdan, J.A.Crowther, G.Otting and N.E.Dixon 2813–2824
- S* The domain structure of *Helicobacter pylori* DnaB helicase: the N-terminal domain can be dispensable for helicase activity whereas the extreme C-terminal region is essential for its function R.G.Nitharwal, S.Paul, A.Dar, N.R.Choudhury, R.K.Soni, D.Prusty, S.Sinha, T.Kashav, G.Mukhopadhyay, T.K.Chaudhuri, S.Gourinath and S.K.Dhar 2861–2874
- Loop II of DNA polymerase beta is important for polymerization activity and fidelity G.C.Lin, J.Jaeger and J.B.Sweasy 2924–2935
- Domain swapping between FEN-1 and XPG defines regions in XPG that mediate nucleotide excision repair activity and substrate specificity M.Hohl, I.Dunand-Sauthier, L.Staresinicic, P.Jaquier-Gubler, F.Thorel, M.Modesti, S.G.Clarkson and O.D.Schärer 3053–3063
- Regulation of B family DNA polymerase fidelity by a conserved active site residue: characterization of M644W, M644L and M644F mutants of yeast DNA polymerase ϵ Z.F.Pursell, I.Isoz, E.-B.Lundström, E.Johansson and T.A.Kunkel 3076–3086
- Nucleoside alpha-thiotriphosphates, polymerases and the exonuclease III analysis of oligonucleotides containing phosphorothioate linkages Z.Yang, A.M.Sismour and S.A.Benner 3118–3127

RNA

- Use of a novel Förster resonance energy transfer method to identify locations of site-bound metal ions in the U2–U6 snRNA complex F.Yuan, L.Griffin, L.Phelps, V.Buschmann, K.Weston and N.L.Greenbaum 2833–2845
- S* Specificity, duplex degradation and subcellular localization of antagomirs J.Krützfeldt, S.Kuwajima, R.Braich, K.G.Rajeev, J.Pena, T.Tuschl, M.Manoharan and M.Stoffel 2885–2892
- S* Thermodynamics and kinetics for base-pair opening in the P1 duplex of the *Tetrahymena* group I ribozyme J.-H.Lee and A.Pardi 2965–2974
- CAF1 plays an important role in mRNA deadenylation separate from its contact to CCR4 T.Ohn, Y.-C.Chiang, D.J.Lee, G.Yao, C.Zhang and C.L.Denis 3002–3015
- S* *Escherichia coli* ribosomal protein L20 binds as a single monomer to its own mRNA bearing two potential binding sites F.Allemand, J.Haentjens, C.Chiaruttini, C.Royer and M.Springer 3016–3031
- S* Base-specific spin-labeling of RNA for structure determination N.Piton, Y.Mu, G.Stock, T.F.Prisner, O.Schiemann and J.W.Engels 3128–3143
- S* Identification of conserved secondary structures and expansion segments in enod40 RNAs reveals new enod40 homologues in plants A.P.Gulyaev and A.Roussis 3144–3152

STRUCTURAL BIOLOGY

- S* Solution structure of Domains IVa and V of the τ subunit of *Escherichia coli* DNA polymerase III and interaction with the α subunit X.-C.Su, S.Jergic, M.A.Keniry, N.E.Dixon and G.Otting 2825–2832
- S* Influence of secondary structure on kinetics and reaction mechanism of DNA hybridization C.Chen, W.Wang, Z.Wang, F.Weil and X.S.Zhao 2875–2884

Continued

<i>S</i>	Solution structure and dynamics of DNA duplexes containing the universal base analogues 5-nitroindole and 5-nitroindole 3-carboxamide	J.Gallego and D.Loakes	2904–2912
<i>S</i>	The open state of human topoisomerase I as probed by molecular dynamics simulation	G.Chillemi, A.Bruselles, P.Fiorani, S.Bueno and A.Desideri	3032–3038
<i>S</i>	Guanines are a quartet's best friend: impact of base substitutions on the kinetics and stability of tetramolecular quadruplexes	J.Gros, F.Rosu, S.Amrane, A.De Cian, V.Gabelica, L.Lacroix and J.-L.Mergny	3064–3075
	The 1.4-Å crystal structure of the <i>S. pombe</i> Pop2p deadenylation subunit unveils the configuration of an active enzyme	A.T.Jonstrup, K.R.Andersen, L.B.Van and D.E.Brodersen	3153–3164
METHODS ONLINE			
	A hybrid CMV-H1 construct improves efficiency of PEI-delivered shRNA in the mouse brain	Z.Hassani, J.-C.François, G.Alfama, G.M.Dubois, M.Paris, C.Giovannangeli and B.A.Demeneix	e65 (9 pp.)
	An <i>in vitro</i> selection scheme for oligonucleotide probes to discriminate between closely related DNA sequences	I.Brukner, R.El-Ramahi, I.Gorska-Flipot, M.Krajcinovic and D.Labuda	e66 (9 pp.)
	Comparison of RNAi efficiency mediated by tetracycline-responsive H1 and U6 promoter variants in mammalian cell lines	J.R.Henriksen, C.Løkke, M.Hammerø, D.Geerts, R.Versteeg, T.Flægstad and C.Einvik	e67 (8 pp.)
	An exonuclease I hydrolysis assay for evaluating G-quadruplex stabilization by small molecules	Y.Yao, Q.Wang, Y.Hao and Z.Tan	e68 (9 pp.)
<i>S</i>	Error-pooling-based statistical methods for identifying novel temporal replication profiles of human chromosomes observed by DNA tiling arrays	T.Park, Y.Kim, S.Bekiranov and J.K.Lee	e69 (14 pp.)
<i>S</i>	Revision of the nonequilibrium thermal dissociation and stringent washing approaches for identification of mixed nucleic acid targets by microarrays	A.E.Pozhitkov, R.D.Stedtfeld, S.A.Hashsham and P.A.Noble	e70 (15 pp.)
	An adaptation of the LMS method to determine expression variations in profiling data	P.Chuchana, D.Marchand, M.Nugoli, C.Rodriguez, N.Molinari and J.A.Garcia-Sanz	e71 (10 pp.)
			3165
CORRIGENDUM			

M, contains a novel method

S, Supplementary Material available at NAR Online